

FAULT CONTROLLED EVOLUTION OF THE VIENNA BASIN

F. MARKO

¹*Department of Geology and Paleontology, Comenius University, Faculty of Sciences, Mlynská dolina, 842 15 Bratislava, Slovakia; marko@nic.fns.uniba.sk*

Abstract: Evolution of the basin has been studied by analysing available field structural data, as well as data compiled from previous mainly published key works as were - map of the preTertiary basement relief, map of the basement units, distribution of sedimentation depocentres, facial character of sediments, subsidence curves and reflexion seismic profiles. Analysing these data active faults pattern was interpreted for each stratigraphic level of the basin, responsible for the shape of sedimentation depocentres. Comparing these synsedimentary faults with fault-slip related paleostress fields, kinematical/dynamical role of faults has been deduced. This approach led to dynamical reconstruction of the basin tectonic evolution.

Key words: fault kinematics, paleostresses, Neogene evolution, Vienna basin

Introduction

The Vienna basin tectonic evolution has complex history due to the exclusive position of the basin in between two different dynamically active Tertiary megablocks - central and outer Western Carpathians, colliding with corner of the Bohemian massif during the formation of the Western Carpathians loop. Complex process of the Western Carpathians propagation to the N and NE prevented by foreland including stress axes as well as rigid body block rotation, fault kinematics fluctuation, fault controlled sedimentation and basin desintegration is recorded in the shape of the basinal sedimentation depocentres and facial type of sedimentary fill. Evolution of the basin has been studied by analysing available field structural data, as well as data compiled from previous mainly published key works as were - map of the preTertiary basement relief, map of the basement units, distribution of sedimentation depocentres, facial character of sediments, subsidence curves and reflexion seismic profiles. Analysing these data active fault pattern was interpreted for each stratigraphic level of the basin, responsible for the shape of sedimentation depocentres.

Comparing these synsedimentary faults with fault-slip related paleostress fields, kinematical/dynamical role of faults has been deduced. This approach led to dynamical reconstruction of the basin tectonic evolution.

Oligocene/Eggenburgian – Early Karpathian (23,4-17 Ma)

First stack of Apulian continental crustal block with NEP happened due to CC collision of northwardly propagated Apulia with Bohemian massif. Terrane in between Apulian promontory and stable Europa plate was extremely shortened, uplifted (Tauren windows, where lowermost crystalline fundament is exhumed). It led to the process of the lateral extrusion combined with gravitational spreading of elastic Carpatho/Pannonian crustal units to the east (Ratschbacher et al. 1991). This so called tectonic escape to the still free space (future Carpatho/Pannonian realm) was controlled by strike-slip faults operating as border faults of crustal wedges moving to the east. During this process already consolidated paleo-mesoalpine structures were within active strike-slip shear zones reworked. The Early Miocene sedimentation of wrench furrows (sensu Montenat et al. 1987) took place within this zones in the regime of dextral transpression (Marko et al. 1991). These basins created E – W trending narrow and long furrows filled up by poorly sorted and shortly transported Eggenburgian – Early Karpathian in age sediments. The Early Miocene wrench furrows were formed at the western part of the internides as well at the eastern part - East Slovakian basin (Vass 1998). To the end of this period due to cca 15° CCW un-block rotation of Carpathians dextral transpression within strike-slips corridors converted to sinistral one and corridors had rotated to ENE - WSW direction. During the Early Miocene period structural evolution was controlled by northward propagation of Apulia promontory and pre-Miocene CCW rotation of Africa plate (Žytko 1982).

Late Karpathian – Lower Badenian (17 – 15,5 Ma)

In between the Early and the Late Karpathian Carpatho/Pannonian units began to move to the north. While Alpine segment of orogene has been already stopped due to collision with Bohemian massif, extruded Carpatho/Pannonian units were pushed to the open space – to the bay of weak crust filled up by Carpathian paleogene flysch sediments. Separation of Carpatho/Pannonian segment from Alpine one and its propagation to the north was realized thanks to N – S dextral strike-slips. One of them is supposed to be in the

basement of the Vienna basin (Krs & Roth 1979, Balla 1988). Within this period started formation of spectacular oroclinal loop. The loop has been traditionally regarded as ductile in nature structure due to bended shape of Alpine units. Others prefer gradual formation of Carpathian loop thanks to independent propagation of detached crustal segments to the Carpathian gulf rimmed by Bohemian massif Tornquist-Teiseire line and Moesia.

During the Late Karpathian ALCAPA invaded to the north, oceanic crust subducted underneath propagating continental Carpatho/Pannonian crustal blocks (Doglioni et al. 1981), was partially melted and produced early volcanic activity of the Western Carpathians.

Distribution of sedimentation depocentres in the Vienna basin really shows dramatical change of structural plan during the Late Karpathian. Structural records show change of compression direction from N - S to NNE - SSW direction, which resulted in change of fault controlled sedimentation kinematics. Former E - W trend of basins was overprinted by NE - SW. NE - SW faults were activated as dominant normal and sinistral strike-slip ones, controlled sedimentation in the Vienna basin and simultaneously allowed northeastward escape of detached blocks after their collision with platforma at the north. During the Late Karpathian first core mountains with exhumed crystalline basement emerged, what is recorded by FT ages (Král 1977, Kováč et al. 1994) as well as by perifault debris sedimentation along uplifted core mountains (Vass et al. 1990). ENE - WSW wrench corridor was active as sinistral transtensional wrench zone. Former Early miocene wrench furrows were broken up and separated remnants were cca 30° - 40° CCW rotated (Kováč & Tunyi 1995). These rotations were coeval with rotations described in Pelso unit (Márton & Fodor 1995).

During the Baddenian due to the activity of major NE - SW sinistral strike-slips pull-apart Vienna basin has opened (Royden et al. 1982, Royden 1985). Baddenian depocentres of sedimentation has N - S trend due to normal faults which accomodated large magnitudes of strike-slip translations. Nevertheless Baddenian Vienna basin is not typical pull-apart, because it has not all attributes of pull-aparts. It was formed as fore-arc basin in relation to the Middle Baddenian volcanic arc. Vienna basin is anomalous from point of mechanism of formation too. Western part of it lies over the accretionary flysch wedge, so it ought to be in position of piggy-back basin, but eastern part lies over the Alpine-Carpathian units of internides.

While transtensional fore-arc basins were formed at the northern edge of the internides, extensional thermal back-arc basins due to lithospheric stretching and volcanism provoked by heat of rising Pannonian astenolith were formed at the Pannonian area.

Middle – Upper Badenian (15,5 – 13,6 Ma)

For this period is typical NE – SW direction of compression and movement of the Carpatho/Pannonian lithospheric blocks. According paleomagnetic data the last CCW block rotations ceased in the western part of Western Carpathians in the Middle Badenian, while the last rotations in the eastern part of the Western Carpathians were recorded at Sarmathian (Orlický 1996). During this period substantial part of volcanic activity had realized, Carpatho-Pannonian volcanic arc was formed. Huge bulk of volcanism was result of melting subducted oceanic/quasioceanic crustal slab, as well as rising of astenospheric astenolith in the Pannonian region (Póka 1988, Lexa et al. 1993). In conditions of thermal crustal extension back arc basins has been formed (Pannonian basin, Danube basin). Crustal stretching due to astenosphere heat was accomodated by extensional normal faulting (Tari et al. 1992, Horváth & Cloetingh 1996).

Sarmathian - Pannonian (11,5 - 9 Ma)

Active front of orogene has already removed far from Western Carpathians area. NE - SW compression gradually inverted to NW - SE pure extension. Former tectonic regime has survived within these conditions. Dominant role as sedimentation controlling structures played NE - SW normal faults. During this period Carpathian lithospheric block propagated to the east. Due to the collision with East European Platform it has stopped movement to the east cca 9 Ma ago. This collision produced E - W compressional event (Decker & Pereson 1996), which affected even such distal terranes as Western Carpathians. E - W normal faults has been reactivated within this stress period and controlled formaion of E - W trending morphostructures - horsts and depressions. Due to steepening of subducting oceanic plate (Royden et al. 1982, Doglioni et al. 1991) and related roll back effect, E - W extension was induced in the Carpathian realm. Numerous N - S faults developed and were reactivated as normal ones within this youngest tensional event. These faults are the most numerous and conspicuous brittle features within the recent architecture of the Western Carpathians, what supports their young age.

Plio - recent (5,3 - 0 Ma)

Pliocene period is typical by lacustrine sedimentation much more modest comparing to former periods of Tertiary sedimentation. Important regional tectonic stresses, generated by

subduction/collision of large lithospheric plates has ceased. Carpathian loop has been already done including Pieniny Klippen Belt structure. There are structural records within Western Carpathians and Pannonian area, that cca N - S (NNW - SSE) compression affected orogen during the end of this period (Csontos et al. 1991, Becker 1993, Hók et al. 1995) and survives within Western Carpathians up till recent as earthquakes focal mechanism analysis of seismogenic faults prove (Pospíšil 1990).

References

- Andrusov, D. 1938: Geologický výskum vnútorného bradlového pásma v Západných Karpatech. III. Tektonika. Rozpr. Stát. geol. Úst. ČSR, 9, 135s.
- Balla, Z. 1987: Tertiary paleomagnetic data for the Carpatho-Pannonian region in the light of Miocene rotation kinematics. *Tectonophysics*, 139, 67-98
- Balla, Z. 1988: A Karpát-Pannon régió nagyszerkezeti képe a felsőeocénben és e kép hatása a mezozoos Tethys-rekonstrukciókra. *Föld. Kozl. Bull. of the Hung. Geol. Soc.*, 118, 11-26
- Birkenmajer, K. 1985: Major strike-slip faults of the Pieniny Klippen Belt and the Tertiary rotation of the Carpathians. *Publ. Inst. Geophys. Pol. Acad. Sci.*, A-16, 175, 101-115
- Csontos, L., Tari, G., Bergerat, F. & Fodor, L. 1991: Evolution of the stress fields in the Carpatho-Pannonian area during the Neogene. *Tectonophysics*, 199, 73-91
- Decker, K. & Pereson, H. 1996: Tertiary kinematics in the Alpine-Carpathian-Pannonian system: links between thrusting, transform faulting and crustal extension. In: Wessely, G. & Liebl, S. (eds.) 1996: Oil and Gas in Alpidic Thrustbelts and Basins of central and Eastern Europe, EAGE Special Publication, 5, 69-77
- Hók, J., Simon, L., Kováč, P., Elečko, M., Vass, D., Verbich, F. & Halmo, J. 1995: Tectonic of the Hornonitrianska kotlina Basin during Tertiary, *Geol. Carpath.*, 46, 191-196
- Horváth, F. & Cloetingh, S. 1996: Stress-induced late-stage subsidence anomalies in the Pannonian basin. *Tectonophysics*, 266, 287-300
- Jiríček, R. 1979: Tektogenetický vývoj Karpatského oblouku během oligocénu a neogénu. In: Mahel', M. (edit.): Tektonické profily Západných Karpát. *Konf., Symp., Semináre, GÚDŠ*, 203-220
- Kováč, M., Baráth, I., Holický, I., Marko, F. & Tunyi, I. 1989: Basin opening in the Lower Miocene strike-slip zone in the SW part of the Western Carpathians. *Geol. Zbor. Geol. Carpath.*, 10, 1, 37-62
- Kováč, M., Král, J., Márton, E., Plašienka, D. & Uher, P., 1994: Alpine uplift history of the Central Western Carpathians: geochronological, paleomagnetic, sedimentary and structural data. *Geol. Carpath.*, 45, 83-96
- Kováč, M. a Tunyi, I. 1995: Interpretácia paleomagnetických údajov zo západnej časti Centrálnych Západných Karpát. *Miner. Slovaca*, 27, 213-220
- Král, J. 1977: Fission track ages of apatites from some granitoid rocks in West Carpathians. *Geol. Zborn. – Geol. Carpath.*, 28, 269-276
- Krs, M. a Roth, Z. 1979: Insubrický karpatský treťohorný blokový systém, jeho vznik a rozloženie. *Geol. zbor., Geol. carpath.*, 30, 1, 3-17
- Lexa, J., Konečný, V., Kaličiak, M. & Hojstričová, V. 1993: Distribúcia vulkanitovkarpato-panónskeho regiónu v priestore a čase. In: Rakús, M. (edit.): Geodynamický model a hlbinná stavba Západných Karpát. *Konf. – symp. – sem., GÚDŠ, Bratislava*, 57-69
- Marko, F., Fodor, L. & Kováč, M. 1991: Miocene strike-slip faulting and block rotation in Brezovské Karpaty Mts. (Western Carpathians). *Min. slovaca*, 23, 189-200
- Márton, E. & Fodor, L. 1995: Combination of paleomagnetic and stress data – a case studies from North Hungary. *Tectonophys.*, 242, 99-114
- Mitchel, A. H. G. & Reading, H. G. 1978: Sedimentation and tectonics. In: Reading, H. G. (edit) *Sedimentary environments and facies*. Elsevier, 439-476
- Montenat, Ch., Ott D'Estvou & Masse, P. 1987: Tectonic-sedimentary characters of the Betic Neogene Basins evolving in a crustal transcurrent shear zone (SE Spain). *Soc. Nat. Elf Aquitaine, Pau, France*, 1-22
- Orlický, O. 1996: Paleomagnetism of neovolcanics of the East-Slovak lowlands and Zemplínske vrchy Mts.: A study of the tectonics applying the paleomagnetic data (W. Carpathians). *Geol. carpathica*, 47/1, 13-20
- Oszczypko, N. 1998: The Western Carpathians Fore-deep development of the foreland basin in front of accretionary wedge and its burial history (Poland). *Geol. carpathica*, 49/6, Bratislava, 415-431
- Póka, T. 1988: Neogene and Quaternary Volcanism of the Carpathian-Pannonian Region: Changes in Chemical composition and its relationship to Basin formation. In: Royden, L. H. & Horváth, F. (eds.): *The Pannonian Basin - a study in basin evolution*. AAPG & Hung. geol. soc., Tulsa, Budapest, 257-278

- Pospíšil, L. 1990: Současné možnosti identifikace střížných zón v oblasti Západních Karpat. *Min. slovac*, 22, 1, 19-31
- Ratschbacher, L., Merle, G., Davy, P. & Cobbold, P. 1991a: Lateral extrusion in the Eastern Alps. Part I: Boundary conditions and experiments scaled for gravity. *Tectonics*, 10, 2, 245-266
- Royden, L. H., Horváth, F. & Burchfiel, B. C. 1982: Transform faulting, extension and subduction in the Carpathian-Pannonian region. *Geol. Soc. An. Bull.*, 93, 715-725
- Royden, L. H. 1985: Vienna Basin a thin skinned pull-apart basin. In: Biddle, K. & Christie Blick, N. (eds.): *SEPM, Spec. publ.*, 37, 319-338
- Tari, G., Horváth, F. & Rumpel, J. 1992: Styles of extension in the Pannonian basin. *Tectonophysics*, 208, 203-216
- Unrug, R. 1984: Geodynamic evolution of the Carpathians. *Ann. Soc. Geol. Pol.*, 52, ¼, 39-66
- Vass, D. 1998: Neogene geodynamic development of the Carpathian arc and associated basins. In: Rakús, M. edit.: *Geodynamic development of the Western Carpathians*. GSSR, Dionýz Štúr Publishers, 155-158
- Vass, D., Nagy, A., Kohút, M. & Kraus, I. 1990: Granitoid clastics on the SE margin of the Vienna basin and basin genesis. In: Minaříková, D. & Lobitzer, H. (eds): *Thirty years of the geological cooperation between Austria and Czechoslovakia*. ÚÚG Praha, 179-184
- Žytko, K. 1982: The Atlantic, the Indian ocean and main linear fracture zones of the post-variscan Europe. *Ann. Soc. Geol. Pol.* 52, ¼, 3-22